arXiv:1208.1993v1 [hep-ph] 9 Aug 2012

On the Gaussian peak of the product of decay probabilities of the standard model Higgs boson at a mass $m_{\rm H} \approx 125$ GeV

David d'Enterria[1]

[1]*CERN, PH Department, 1211 Geneva, Switzerland*

The product of the branching ratios of the standard model (SM) Higgs boson in all decay channels available below the top-antitop threshold is observed to be a Gaussian distribution of the Higgs boson mass with a maximum centred around $m_{\rm H} \approx 125$ GeV, i.e. in the region of masses where a new boson has been discovered at the Large Hadron Collider. This fact places the Higgs particle at a mass of “maximum opportunity’ in terms of the study of its decays and couplings to other particles. Such an intriguing observation is seemingly driven by the different $m_{\rm H}$-power dependence of the Higgs decay-widths into gauge bosons and fermions featuring a steep anticorrelated evolution below the WW decay threshold in the transition region of masses $m_{\rm H} \approx m_{\rm W} - 2m_{\rm W}$. Speculative consequences of taking this seemingly accidental feature as indicative of an (unknown) underlying dynamics of the SM are also discussed.

PACS numbers: 14.80.Bn

I. INTRODUCTION

In the standard model (SM) of particle physics, the masses of the electroweak W and Z vector bosons are generated via the Higgs-Brout-Englert-Guralnik-Hagen-Kibble mechanism [1] which also predicts the existence of a new elementary neutral scalar (Higgs) boson. The fermion masses are also generated through Yukawa interactions with the Higgs field. The mass of the Higgs boson itself, given by the square-root of its (unknown) self-coupling parameter λ ($m_{\rm H}^2 = \sqrt{2}\lambda/G_{\rm F}$ with Fermi constant $G_{\rm F} \approx 1.16637\cdot10^{-5}$ GeV^{-2}) is however a free parameter of the theory. Theoretically, the value of $m_{\rm H}$ is constrained from above by requirements of unitarity of the elastic $\rm W_L W_L$ cross sections [2] as well as by triviality bounds for large self-couplings λ [3]. Lower limits on $m_{\rm H}$ are imposed by vacuum stability arguments (λ cannot be too small since otherwise the top-quark Yukawa coupling would make it negative) [3]. The recent experimental observation at the Large Hadron Collider (LHC) of a new boson at $m_{\rm H} \approx 125$ GeV in various decay modes consistent with the hypothesized SM Higgs boson [4, 5], constitutes a experimental breakthrough in our understanding of the mechanism by which all elementary particles acquire their mass.

Basic conservation rules imply that a neutral spinless SM Higgs boson can only decay into (i) pairs of fundamental fermions $\rm q\bar{q} = u\bar{u}, d\bar{d}, s\bar{s}, c\bar{c}, b\bar{b}, t\bar{t}$ for quarks and $\ell^+\ell^- = \rm e^+e^-, \mu^+\mu^-, \tau^+\tau^-$ for leptons (neglecting neutrinos which are massless in the SM, see below), (ii) pairs of (virtual or real) heavy bosons VV = ZZ,WW; as well as (iii) pairs of massless bosons ($gg, \gamma\gamma$, as well as $Z\gamma$) via heavy-particle (mostly W and top-quark) loops. The partial decays widths Γ_i of the Higgs boson into the most relevant channels, i.e. the inverse decay rates to each final state kinematically allowed for a given value of $m_{\rm H}$, have been determined including various high-order quantum chromodynamics (QCD) and electroweak corrections (see [6] and refs. therein). The total width is obtained by summing all the Higgs partial widths: $\Gamma_{\rm tot} = \sum_i \Gamma_i$. The branching ratio of a given mode is $\mathrm{BR} = \Gamma_i/\Gamma_{\rm tot}$ and, of course, the sum of all of them amounts to one, $\sum_i \mathrm{BR}_i = 1$. Decay widths and branching ratios to the various final-states can be computed using programs such as HDECAY [6], and have typical theoretical uncertainties in the range $\Delta\mathrm{BR} \approx 3$–10% for $m_{\rm H} \approx 125$ GeV [7]. The SM Higgs couplings to fundamental fermions are proportional to the fermion masses $m_{\rm f}$, whereas its couplings to bosons are proportional to the square of their masses $m_{\rm V}^2$, a clear indication of the primary role of the Higgs particle in the breaking of electroweak symmetry. The partial $\rm f\bar{f}$ and VV decay-widths are thus given by the tree-level relations (see e.g. [8])

$$\Gamma_{\rm f\bar{f}} = \frac{N_c\, G_{\rm F}\, m_{\rm f}^2}{4\sqrt{2}\pi} \cdot m_{\rm H} \cdot \left(1 - \frac{4m_{\rm f}^2}{m_{\rm H}^2}\right)^{3/2}, \quad \text{where } N_c = 3\,(1) \text{ for quarks (leptons)}, \tag{1}$$

$$\Gamma_{\rm VV} = \frac{\delta_V\, G_{\rm F}}{16\sqrt{2}\pi} \cdot m_{\rm H}^3 \cdot \sqrt{1 - \frac{4m_{\rm V}^2}{m_{\rm H}^2}} \left(1 - \frac{4m_{\rm V}^2}{m_{\rm H}^2} + \frac{3}{4}\left(\frac{4m_{\rm V}^2}{m_{\rm H}^2}\right)^2\right), \quad \text{with } \delta_{\rm V} = 2\,(1) \text{ for W (Z)}. \tag{2}$$

From these expressions one sees first that the decays to fermions are basically linear in $m_{\rm H}$ but follow a $m_{\rm H}^3$ dependence for bosons, i.e. for large enough $m_{\rm H}$ the decays to W and Z are clearly preferred over fermions. [The loop-mediated gg and $\gamma\,\gamma$, Z γ decay widths – with more complex expressions not reproduced here – feature also a cubic dependence on $m_{\rm H}$, but with a smaller $\Gamma_{\rm loop}$-prefactor than for VV decays]. Equation (2) is only valid for on-shell vector bosons, i.e. for $m_{\rm H} \geq 2m_{\rm V}$. In the region where one or both W and Z particles can be produced off-shell, the corresponding partial widths are smaller but rise much faster with the Higgs mass (a power-law fit to $\Gamma_{\rm V^*V}(m_{\rm H})$ in the range $m_{\rm H} \approx m_{\rm Z}$–$2m_{\rm W}$ computed with HDECAY shows a dependence as fast as $m_{\rm H}^{14}$). As result, the subtreshold WW* and ZZ* branching ratios are larger than those for fermions already for $m_{\rm H} \approx 135$ GeV and 157 GeV respectively (Fig. 1).

In this work, we investigate whether there is a particular mass region for which the SM Higgs boson has the maximum number of decay modes effectively available. More precisely, we want to see which is the value (if any) of $m_{\rm H}$ which maximizes the distribution of the product of branching ratios

$$\mathrm{BR}_{\rm prod}(m_{\rm H}) \; = \; \prod_i \, \mathrm{BR}_{\rm H\to i}(m_{\rm H})\,. \tag{3}$$

One finds somehow surprisingly that the distribution $\mathrm{BR}_{\rm prod}(m_{\rm H})$ is Gaussian in the range of masses $m_{\rm H} \approx m_{\rm Z}$–$2m_{\rm W}$ with a maximum centred around $m_{\rm H} \approx 125$ GeV and a width of $\sigma_{m_{\rm H}} \approx 6$ GeV. The reasons of such a behaviour are explored below.

II. THEORETICAL SETUP

The partial widths and branching ratios of the Higgs boson in the SM are determined following the same approach developed in Refs. [7, 9]. We use HDECAY (version 4.43) [6] for the fermionic, digluon, diphoton and Zγ decays, and PROPHECY4F (version 2.0) [10] for the weak boson decays. HDECAY includes next-to-leading order (NLO) corrections (or higher-order ones in some cases) for the QCD and electroweak contributions for all decay channels, whereas PROPHECY4F takes into account complete NLO and electroweak corrections as well as interferences for all final-states in the W,Z decays. Both calculations are then combined by modifying the Higgs total width obtained with HDECAY according to $\Gamma_{\rm H} \;=\; \Gamma^{\rm HD}_{\rm H} - \Gamma^{\rm HD}_{\rm ZZ} - \Gamma^{\rm HD}_{\rm WW} + \Gamma^{\rm Proph}_{\rm 4f}$ [7], where $\Gamma^{\rm HD}$, $\Gamma^{\rm HD}_{\rm ZZ}$ and $\Gamma^{\rm HD}_{\rm WW}$ stand respectively for the total, and partial widths to ZZ and WW calculated with HDECAY, and $\Gamma^{\rm Proph}_{\rm 4f}$ represents the partial width of H → 4f, including interferences, calculated with PROPHECY4F. The results of PROPHECY4F modify by about ±10% the branching ratios computed with HDECAY alone mostly in the mass region $m_{\rm H} = m_{\rm Z}$–$2m_{\rm W}$. For the input parameters of the calculations the latest PDG values [11] are employed (Table I), with the uncertainties quoted for a few parameters being the same as those used in [7, 9]. The strange-quark mass used is the $\overline{\rm MS}$ value at two different scales $Q = 1, 2$ GeV, and the charm and bottom quark masses correspond to the 1-loop pole values which are less sensitive to α_s uncertainties.

TABLE I: Input parameters used for the Higgs boson branching ratios (BR) calculations carried out in this work.

$m_{\rm s}$ (GeV)	0.1(2 GeV),0.19(1 GeV)	$m_{\rm e}$ (MeV)	0.510998928	$m_{\rm W}$ (GeV)	80.385	$\alpha(0)$	1/137.035999074
$m_{\rm c}$ (GeV)	1.28 ± 0.03	m_μ (MeV)	105.658367	$m_{\rm Z}$ (GeV)	91.1876	$G_{\rm F}$ (GeV^{-2})	$1.1663787 \cdot 10^{-5}$
$m_{\rm b}$ (GeV)	4.16 ± 0.06	m_τ (MeV)	1776.82	$\Gamma_{\rm W}$ (GeV)	2.085	$\alpha_{\rm s}(m_{\rm Z}^2)$	0.1184 ± 0.002
$m_{\rm t}$ (GeV)	173.5 ± 2.5			$\Gamma_{\rm Z}$ (GeV)	2.4952		

The Higgs boson decays into the first-generation fermions ($\mathrm{u\,\overline{u}}$, $\mathrm{d\overline{d}}$, $\mathrm{e^+e^-}$) are not directly implemented in HDECAY since, due to their small masses compared to $m_{\rm H}$ they have extremely reduced branching ratios and are of no experimental relevance. From Eq. (1) the dependence of the first-family branching-fractions on $m_{\rm H}$ can be obtained from the corresponding widths for heavier (e.g. second-generation) fermions via

$$\mathrm{BR}_{\rm u\overline{u},d\overline{d}}(m_{\rm H}) \;\approx\; \left(\frac{m_{\rm u,d}}{m_{\rm s}}\right)^2 \cdot \mathrm{BR}_{\rm s\overline{s}}(m_{\rm H})\,, \quad \mathrm{BR}_{\rm e^+e^-}(m_{\rm H}) \;\approx\; \left(\frac{m_{\rm e^+e^-}}{m_{\mu^+\mu^-}}\right)^2 \cdot \mathrm{BR}_{\mu^+\mu^-}(m_{\rm H})\,. \tag{4}$$

The branching ratios for the quarks of the first generation are obtained using this expression with $m_{\rm u,d,s} \approx 3, 5, 100$ MeV. Although the resulting $\mathrm{u\,\overline{u}}$, $\mathrm{d\overline{d}}$ ($\mathrm{e^+e^-}$) decays are suppressed by factors of order 10^{-3} (10^{-5}) compared to the $\mathrm{s\,\overline{s}}$ ($\mu^+\mu^-$) channel, the dependence of their BR on $m_{\rm H}$ is the same as for other heavier fermions.

Since the light-quark masses are not precisely known, there are potentially large uncertainties in their associated BR values. As a cross-check we computed the branching ratios using the two values for the s-quark mass quoted in Table I and found that although the corresponding absolute $\mathrm{BR}_{u\bar{u},d\bar{d},s\bar{s}}$ values changed by a factor of two, the m_H-dependence of $\mathrm{BR}_{u\bar{u},d\bar{d},s\bar{s}}(m_H)$ is the same for any Higgs mass value. Similarly, the same changes of $m_{u,d,s}$ modify the total Higgs decay width only at the permille level given the smallness of these suppressed branching ratios. Since we are basically interested in the *shapes* of $\mathrm{BR}(m_H)$, and since unc ertainties in $\mathrm{BR}_{u,d,s}$ propagate just as an overall normalization factor of $\mathrm{BR}_{\mathrm{prod}}(m_H)$, we do not take into account the u,d,s mass uncertainties hereafter.

III. RESULTS

The branching ratios for all kinematically allowed channels of the SM Higgs boson are computed for masses $m_H \approx 0$–1 TeV using the setup discussed in the previous Section. Figure 1 shows the nine fermionic, two bosonic, and three loop-mediated BR in the mass range $m_H = 10$–1000 GeV. Several observations are worth pointing out. The eight lightest fermionic channels (plotted here well above their respective thresholds at $m_H = 2m_{f\bar{f}}$) have a relatively flat BR as a function of m_H up to about $\simeq$110 GeV, where the very fast-rising VV* decays start to play an increasing role which subsequently make all the $f\bar{f}$ branching fractions drop equally steeply. Beyond $2m_V$ the bosonic decays are roughly constant as a function of m_H except for the relatively small modulation due to the onset of the $t\bar{t}$ mode around $m_H \approx 2m_t$, while all other fermionic decays decrease monotonically. The loop-mediated bosonic channels ($gg, \gamma\gamma, Z\gamma$) show first a rise with m_H peaking at $m_H \approx 120$–140 GeV followed also by a monotonic falloff. It is



FIG. 1: Decay branching fractions of the Higgs boson into all fermion and boson pairs of the SM, as a function of the Higgs boson mass, computed with HDECAY and PROPHECY4F with the parameters listed in Table I and using Eq. (4) for the first-generation fermions. The dashed vertical line indicates the position of the 125-GeV boson resonance observed at the LHC [4, 5].

clear from Fig. 1 that all the "action" occurs over the mass region (m_W, $2m_W$) centered at around $m_H \approx 125$ GeV (dashed line in Fig. 1). The value of the W mass and the preferred decay of the SM Higgs boson into weak bosons are responsible for all the features seen in the distribution of branching ratios over this distinctive transition mass region. Indeed, the fast rise of the bosonic decays, the consequent fast drop of all (light) fermionic modes, and the maxima of the loop-mediated decays around the region of the newly observed LHC boson resonance, are all just driven by the particular value of the W boson mass[1].

Figure 2 shows the product of all computed branching ratios as a function of $m_{\rm H}$, Eq. (3), below the $t\bar{t}$ threshold[2] plotted in logarithmic (left) and linear (right) scales. The band/boxes around the curves indicate the corresponding theoretical and parametric uncertainties, estimated as discussed in [7], propagating only those which are uncorrelated across the individual decay channels. The absolute amplitude of $\mathrm{BR}_{\rm prod}(m_{\rm H})$ has no particular meaning and the distribution has been normalized by its integral. We note that below m_Z, the $Z\gamma$-decay is actually null in HDECAY since the code does not include the $Z^*\gamma$ channel[3]. This fact explains the sudden drop seen in $\mathrm{BR}_{\rm prod}(m_{\rm H})$ around 90 GeV. The resulting distribution can be very well fitted by a Gaussian function in the transition region from the fermion- to the boson-dominated regime $m_{\rm H} = m_Z$–$2m_W$ with goodness-of-fit per degree-of-freedom: $\chi^2/\mathrm{ndf} = 61.6/68$. Surprisingly, the distribution has not only a Gaussian shape but it has a maximum centered at $m_{\rm H} \approx 125$ GeV (and a width of about $\sigma_{m_{\rm H}} = 6$ GeV) i.e. in the range of masses where the Higgs-like particle has been discovered at the LHC: $m_{\rm H}^{\rm CMS} = 125.3 \pm 0.6$ [4] and $m_{\rm H}^{\rm ATLAS} = 126.0 \pm 0.6$ [5] (hatched grey band in Fig. 2). Although the fit quality ($\chi^2/\mathrm{ndf} \approx 1$) is very good within $3\sigma_{m_{\rm H}}$, the product of branching ratios starts to deviate from a Gaussian in the tail beyond $m_{\rm H} \approx 155$ GeV. This disagreement may be corrected if one adds the $t\bar{t}$ Higgs decays including the 4-body contributions from diagrams involving two virtual top-quarks which are very small in amplitude (although steeply rising with $m_{\rm H}$) and currently neglected in HDECAY.



FIG. 2: Product of all decay branching ratios of the SM Higgs particle into fermion and boson pairs below the $t\bar{t}$ threshold as a function of the Higgs boson mass, Eq. (3) normalized to its integral. The distribution is shown in log (left) and linear (right) scales and fitted to a Gaussian with the parameters listed in the legends. The uncorrelated uncertainties obtained in [7] are shown as boxes/band. The hatched vertical band indicates the position of the observed boson resonance at the LHC [4, 5].

We have also checked what happens with $\mathrm{BR}_{\rm prod}(m_{\rm H})$ if one excludes one or various decay channels. As a matter of fact, the product of the two dominant fermion ($b\bar{b}$) and bosonic (WW) decay branching ratios is already a distribution with Gaussian-like shape and a maximum at $m_{\mathrm{H}} \approx 135$ GeV with larger width $\sigma_{m_{\mathrm{H}}} \approx 14$ GeV. The product of the three loop-mediated branching fractions alone is also normally distributed around $m_{\mathrm{H}} \approx 131$ GeV with a width $\sigma_{m_{\mathrm{H}}} \approx 11$ GeV. The consecutive addition of extra fermion-pair BR displaces the Gaussian to lower m_{H} values and reduces its associated width, resulting in the final overall distribution plotted in Fig. 2.

---

[1] In the SM, the Z and W masses are linked through the Weinberg mixing angle.

[2] The $t\bar{t}$ branching ratio is actually set to zero in HDECAY below $m_{\rm H} \approx 250$ GeV and is omitted in the product of $\mathrm{BR}(m_{\rm H})$ distributions.

[3] Likewise, HDECAY does not include the Dalitz $\gamma^*\gamma$ mode with γ^* decaying into pairs of fermions – amounting to about 10% of the $\gamma\gamma$ decay ratio according to [12] – which nonetheless has the same $m_{\rm H}$-dependence as $\mathrm{BR}_{\gamma\gamma}$ and thus does not affect the shape of the obtained $\mathrm{BR}_{\rm prod}(m_{\rm H})$ distribution.

The first pragmatical consequence of the observation plotted in Fig. 2 is that a SM Higgs boson around 125 GeV has the largest possible number of decay modes concurrently open to potential experimental study. No other Higgs mass value has theoretically a better "signal strength" for the *whole* set of kinematically-allowed decay channels simultaneously. This fact places the Higgs boson in a mass window of "maximum opportunity" to experimentally study its couplings to all the gauge bosons and fermions at the LHC as well as at any other future collider.

A priori, there is no physical reason why the product of the many (or a few) branching-ratios distributions plotted in Fig. 1 should be normally distributed. However, in statistics theory it is well-known that probability densities such as the beta distribution, $P(x) \propto x^m(1-x)^n$ for $0 < x < 1$, and the gamma distribution, $P(x) \propto x^m \exp(-x)$ for $x > 0$, converge to Gaussian distributions[4] for large values of the powers m and n. From the tree-level Higgs-mass dependencies of the decay widths given by Eqs. (1) and (2): $\Gamma_{\mathrm{f\bar{f}}} \propto m_{\mathrm{H}}$, $\Gamma_{\mathrm{VV}} \propto m_{\mathrm{H}}^3$ (and $\Gamma_{\mathrm{loop}} \propto m_{\mathrm{H}}^3$), the product of branching ratios for the fermionic and bosonic (and loop-mediated) decays is indeed a expression which has the following generic parametric form

$$\mathrm{BR}_{\mathrm{prod}}(m_{\mathrm{H}}) \;\propto\; \left(\prod_i^n \Gamma_i(m_{\mathrm{H}})\right) \times \left(\sum_i^n \Gamma_i(m_{\mathrm{H}})\right)^{-n} \propto \left(m_{\mathrm{H}}^{n_{\mathrm{f\bar{f}}}} \cdot m_{\mathrm{H}}^{3(n_{\mathrm{VV}}+n_{\mathrm{loop}})}\right) \times \left(A \cdot n_{\mathrm{f\bar{f}}} \cdot m_{\mathrm{H}} + B \cdot (n_{\mathrm{VV}} + n_{\mathrm{loop}}) \cdot m_{\mathrm{H}}^3\right)^{-n} ,$$

for a total number of decay modes $n = n_{\mathrm{f\bar{f}}} + n_{\mathrm{VV}} + n_{\mathrm{loop}} = 13$ (below the $\mathrm{t\bar{t}}$ threshold) and constants A and B. Taking $n_{\mathrm{f\bar{f}}} = 8$, $n_{\mathrm{VV}} = 2$ and $n_{\mathrm{loop}} = 3$ one obtains a beta-type distribution with large exponents

$$\mathrm{BR}_{\mathrm{prod}}(m_{\mathrm{H}}) \;\propto\; m_{\mathrm{H}}^{10} \times \left(A' + B' \cdot m_{\mathrm{H}}^2\right)^{-13} \tag{5}$$

which can be indeed a Gaussian-like function for appropriate positive A' and B' constants.

Although the observed Gaussian shape of $\mathrm{BR}_{\mathrm{prod}}(m_{\mathrm{H}})$ seems to be accidental and it happens to peak around $m_{\mathrm{H}} \approx 1.5 m_{\mathrm{W}}$ for the reasons described above, one can speculate about the possibility that there could be an (unknown) underlying dynamical reason that makes the product of SM Higgs boson branching fractions to peak at m_{H} and, therefore, use this fact to place limits or constraints on new possible particles and/or Higgs decay channels. Making a simple analogy with entropy arguments in the decay of an excited system in equilibrium, one could naively argue that given that the physical realization of the Higgs field – which gives mass to the rest of elementary particles – is an unstable particle, the value of m_{H} must be such that the Higgs boson has the largest number of ways in which its available mass can be potentially distributed into the known fundamental particles. Thus, if the 125-GeV resonance observed at the LHC is indeed the SM Higgs boson and the maximum of the product of its branching-ratios at m_{H} derives from some more fundamental SM ground one could constrain the Higgs "invisible" decays into e.g. neutrinos or axions [13]. Adding, for example, three new Higgs boson decays into (Dirac) neutrino-antineutrino pairs with Yukawa couplings as for the other SM fermions, i.e. via Eq. (4) using m_ν instead of m_e, displaces the peak of the distribution to about $m_{\mathrm{H}} \approx 122$ GeV. It would be interesting to test the shape of the corresponding $\mathrm{BR}_{\mathrm{prod}}(m_{\mathrm{H}})$ distribution taking into account different neutrino mass realizations e.g. in the context of seesaw mechanisms. Similarly, we defer for a coming study to check whether the m_{H}-peak in the product of decay fractions present in the SM appears also for the (unconstrained) mass of other Higgs particles in the context of supersymmetric (SUSY), or more generally two-Higgs-doublet, extensions of the SM [14].

IV. SUMMARY

The fourteen branching ratios of the standard model Higgs boson into the known fermion and boson pairs have been computed as a function of the Higgs boson mass in the range $m_{\mathrm{H}} \approx 0$–1 TeV with the HDECAY and PROPHECY4F programs. The product of the branching ratios in all decay channels available below the top-antitop threshold is observed to be a Gaussian distribution of the Higgs boson mass with a maximum centred around $m_{\mathrm{H}} \approx 125$ GeV, i.e. in the region of masses where a new boson resonance has been discovered at the LHC. We have argued that such an intriguing feature is likely the accidental result of the fact that the fermionic[5], bosonic and loop-mediated Higgs decay branching ratios feature all strongly anti-correlated changes around $m_{\mathrm{H}} \approx 1.5 m_{\mathrm{W}}$ due to their different m_{H}-power dependencies. As a result, the product of their branching ratios has a beta-type probability distribution which converges into a Gaussian for large number of decay channels. If, on the contrary, this observation is due to an (unknown) underlying physical mechanism – i.e. if there is any fundamental reason why the value of m_{H} must be such that it has the largest number of ways in which the available mass can be potentially distributed into the known fundamental particles – such an effect could be used to impose extra constraints on new possible particles and/or Higgs decay channels in the SM and its extensions.

---

[4] With means $\mu = m/(m+n)$, m and widths $\sigma = (m \cdot n)/(m+n)^3$, m for the beta and gamma distributions respectively.

In any case, the observation reported here places the Higgs boson in a mass window of "maximum opportunity" to experimentally study its couplings to all the gauge bosons and fermions, and therefore impose maximum constraints to any physics beyond the standard Model, at the LHC as well as at any other future collider.

Acknowledgments

Valuable discussions with Andre David, Albert de Roeck and Michelangelo Mangano are acknowledged. I gratefully acknowledge partial support of the Perimeter Institute for Theoretical Physics and of Brookhaven National Lab where part of this work was undertaken.

[1] F. Englert and R. Brout, Phys. Rev. Lett. **13**, 321 (1964); P. W. Higgs, Phys. Lett. **12** (1964) 132; P. W. Higgs, Phys. Rev. Lett. **13**, 508 (1964); G. S. Guralnik, C. R. Hagen and T. W. B. Kibble, Phys. Rev. Lett. **13** (1964) 585; P. W. Higgs, Phys. Rev. **145** (1966) 1156; T. W. B. Kibble, Phys. Rev. **155** (1967) 1554.

[2] M. J. G. Veltman, Acta Phys. Polon. B **8** (1977) 475; B. W. Lee, C. Quigg and H. B. Thacker, Phys. Rev. Lett. **38** (1977) 883; Phys. Rev. D **16** (1977) 1519.

[3] N. Cabibbo, L. Maiani, G. Parisi and R. Petronzio, Nucl. Phys. B **158** (1979) 295.

[4] S. Chatrchyan *et al.* [CMS Collaboration], arXiv:1207.7235 [hep-ex].

[5] G. Aad *et al.* [ATLAS Collaboration], arXiv:1207.7214 [hep-ex].

[6] M. Spira, Nucl. Instrum. Meth. A **389** (1997) 357; A. Djouadi, J. Kalinowski and M. Spira, Comput. Phys. Commun. **108** (1998) 56; `http://people.web.psi.ch/spira/hdecay/`.

[7] A. Denner, S. Heinemeyer, I. Puljak, D. Rebuzzi and M. Spira, Eur. Phys. J. C **71** (2011) 1753.

[8] A. Djouadi, Phys. Rept. **457**, 1 (2008).

[9] S. Dittmaier, S. Dittmaier, C. Mariotti, G. Passarino, R. Tanaka, S. Alekhin, J. Alwall and E. A. Bagnaschi *et al.*, arXiv:1201.3084 [hep-ph].

[10] A. Bredenstein, A. Denner, S. Dittmaier and S. Pozzorini, Phys. Rev. Lett. **103** (2009) 012002; JHEP **0808** (2008) 108; JHEP **1003** (2010) 021; `http://omnibus.uni-freiburg.de/~sd565/programs/prophecy4f/prophecy4f.html`

[11] J. Beringer *et al.* [Particle Data Group], Phys. Rev. D86 (2012) 010001.

[12] A. Firan and R. Stroynowski, Phys. Rev. D **76** (2007) 057301.

[13] R. E. Shrock and M. Suzuki, Phys. Lett. B **110** (1982) 250.

[14] D. d'Enterria, in preparation.

---

[5] Note that this statement applies to all fermion pairs, including the top-quark which naively has a much larger scale ($2m_{\mathrm{t}}$) than m_{W}. Indeed, the shape of the $\mathrm{t\bar{t}}$ branching ratio also directly depends on the value of m_{W} in the region where one or both top-quarks are off-shell. Also, the top-quark together with the W boson enter indirectly into the loop-mediated decays involving massless gauge bosons and determine their behaviour (local maxima) in the region $m_{\mathrm{H}} = (m_{\mathrm{W}}, 2m_{\mathrm{W}})$.